

# JACOBIANS

If  $u$  and  $v$  are functions of two independent variables  $x$  and  $y$ , then the determinant  $\begin{vmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{vmatrix}$  is called Jacobian of  $u, v$  and denoted by  $J(u, v) = \frac{\partial(u, v)}{\partial(x, y)}$

Similarly

$$J(u, v, w) = \frac{\partial(u, v, w)}{\partial(x, y, z)} = \begin{vmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} & \frac{\partial u}{\partial z} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} & \frac{\partial v}{\partial z} \\ \frac{\partial w}{\partial x} & \frac{\partial w}{\partial y} & \frac{\partial w}{\partial z} \end{vmatrix}$$

## Properties of Jacobians

### (1) Chain rule for Jacobians

If  $u(x, y)$  and  $v(x, y)$  i.e.  $u$  and  $v$  are fun. of  $x, y$  and  $x, y$  are fun. of  $(r, s)$  then

$$\frac{\partial(u, v)}{\partial(x, y)} = \frac{\partial(u, v)}{\partial(r, s)} \times \frac{\partial(r, s)}{\partial(x, y)}$$

(2) If  $J_1$  is Jacobian of  $(u, v)$  wr.t.  $(x, y)$  and  $J_2$  is the Jacobian of  $(x, y)$  wr.t.  $(u, v)$  then  $J_1 J_2 = 1$  i.e.

$$\frac{\partial(u, v)}{\partial(x, y)} \times \frac{\partial(x, y)}{\partial(u, v)} = 1$$

Ex 1. If  $u = x(1-y)$ ,  $v = xy$  Prove that  $JJ' = 1$ .

2. If  $x = r \cos \theta$ ,  $y = r \sin \theta$  Show that  $\frac{\partial(x, y)}{\partial(r, \theta)} = r$  and  $JJ' = 1$

3. If  $y_1 = \frac{x_2 x_3}{x_1}$ ,  $y_2 = \frac{x_3 x_1}{x_2}$ ,  $y_3 = \frac{x_1 x_2}{x_3}$ , show that

$$\frac{\partial(y_1, y_2, y_3)}{\partial(x_1, x_2, x_3)} = 4$$